

## Global Analysis of Estrogen-Regulated Genes in Mouse Uterus using cDNA Microarray and Laser Capture Microdissection

Seok Ho Hong<sup>1,2</sup>, Hee Young Nah<sup>2</sup>, Ji Yoon Lee<sup>2</sup>,  
Chung Hoon Kim<sup>2</sup>, Moon Kyo Kim<sup>1</sup>

<sup>1</sup>Department of Life Science, College of Natural Sciences, Hanyang University, Seoul,  
<sup>2</sup>Department of Obstetrics and Gynecology, College of Medicine, Ulsan University,  
Asan Medical Center, Seoul, Korea

### cDNA Microarray와 Laser Capture Microdissection을 이용한 생쥐 자궁에서 Estrogen에 의해 조절되는 유전자 발현에 관한 분석

한양대학교 자연과학대학 생명과학과<sup>1</sup>, 울산대학교 의과대학 서울아산병원 산부인과<sup>2</sup>

홍석호<sup>1,2</sup>, 나희영<sup>2</sup>, 이지윤<sup>2</sup>, 김정훈<sup>2</sup>, 김문규<sup>1</sup>

연구목적: Estrogen

estrogen

연구재료 및 방법: 8, 14, estrogen, cDNA microarray, Estrogen, 6, 12, laser capture microdissection (LCM), estrogen

결과: Estrogen 6, 가 0.9% (가 22, 49), 12, 8.4% (가 351, 287), 가, 가

LCM

결론: estrogen, 가, 1) estrogen, 12, , 2), 가

**Key Words:** Estrogen, Uterus, cDNA microarray, Laser capture microdissection

The steroid hormones, estrogen and progesterone, play a pivotal role in the development of reproductive tract and regulation of implantation process. The cellular actions of these steroid hor-

: , ) 133-791, 17,  
Tel: (02) 2290-0954, Fax: (02) 2290-1960, e-mail: kimmk@email.hanyang.ac.kr  
: , ) 138-736, 388-1,  
Tel: (02) 3010-5053, Fax: (02) 3010-5059, e-mail: shhong@amc.seoul.kr

mones are mediated through their ability to bind nuclear receptors, which are basically ligand-activated transcription factors. Estrogen receptors (ERs) are known to exist at least two types, ER $\alpha$  and ER $\beta$ , in mammals.<sup>1,2</sup> Estrogen-ER complex directly binds to estrogen response elements on target genes and modulates gene transcription in a ligand-dependent manner, so-called genomic actions of estrogen.<sup>3</sup> However, some estrogen actions are mediated by intracellular second messengers or various signal-transduction cascades through the mechanism of nongenomic actions.<sup>4,5</sup> Together with the brain, the uterus is a major estrogen target organ, which is composed of heterogeneous cell types that respond differentially to estrogen. To date, although relatively few genes have been identified in the uterus that are regulated by estrogen, the exact molecular pathways of estrogen and the expression patterns of specific cell types are largely unknown.<sup>6-10</sup>

The aim of this study was to identify early- and late-responsive genes regulated by estrogen in the mouse uterus and determine whether estrogen-responsive genes are differentially expressed according to the uterine specific cell types. The technique of cDNA microarray has been used to profile the genome-wide analysis of gene expression in various uterine tissues such as implantation-interimplantation sites, progesterone-induced and delayed-implanted uterus.<sup>11,12</sup> We also employed a combination of laser capture microdissection (LCM) and RT-PCR to quantify estrogen-responsive genes in specific cell types of the ovariectomized mouse uterus. In recent, LCM was also used to isolate of specific cell types from tissue sections and examine the differentially expressed genes.<sup>13</sup> Using these technique, we profiled both early- and late-responsive genes regulated by estrogen and demonstrated that several responsive genes was differentially expressed in specific cell types of the estrogen-induced uterus.

## MATERIALS AND METHODS

### 1. Animals and Estrogen treatments

ICR mice were housed within temperature- and light-controlled conditions under the supervision of a licenced veterinarian. Mice were maintained on a 12L : 12D photoperiod and provided with food and water ad libitum. Female mice (6~7 weeks of age) were ovariectomized (OVX) and rested for 14 days before receiving estrogen treatment. They were injected with oil (0.1 ml) and 17 $\beta$ -estradiol (Sigma, St. Louis, MO; 300 ng/mouse) which was dissolved in sesame oil and injected (0.1 ml/mouse) subcutaneously. Mice were killed and uterine horns (n=9) were collected at 6 h or 12 h after injection. All animal experiments were performed in accordance with the Guide of Ulsan University for Care and Use of Laboratory Animals.

### 2. RNA isolation for cDNA microarray

Uterine tissues from nine female mice were pooled, snap frozen and homogenized by mortar in liquid nitrogen. For cDNA microarray analysis, total RNA was extracted using TRIZOL reagent (InVitrogen, Carlsbad, CA) and purified using RNeasy total RNA isolation kit (Qiagen, Valencia, CA) according to the manufacturers instructions. Total RNA was quantified by spectrophotometer and its integrity was assessed by running on a denaturing 0.8% agarose gel. Prior to use in cDNA microarray analysis, each RNA sample via OVX/estrogen treatment/6 h/12 h protocol was validated by assaying for up-regulation (Asparagine synthetase and Lactoferrin) or down-regulation genes (Glutathione S-transferase and Secreted frizzled-related sequence protein 2) as markers of estrogen efficacy.

### 3. cDNA microarray and Data analysis

Profiling of estrogen-regulated gene expression

was analyzed with a TwinChip Mouse-7.4 K (Digital Genomics, Seoul, Korea) consisting of 7616 mouse cDNA clones. Twenty  $\mu\text{g}$  RNA was reverse-transcribed with Cy3- or Cy5-conjugated dUTP (Amersham Pharmacia Biotech, Piscataway, NJ) respectively, using SuperScript II (Gibco BRL, Rockville, MD) and oligo(dT)<sub>18</sub> primer (Ambion, Austin, TX) in a reaction volume of 20  $\mu\text{l}$  according to the method suggested by the manufacturer. After the labeling reaction for 1 h at 42 °C, unincorporated fluorescent nucleotide was cleaned up using Microcon YM-30 column (Millipore, Bedford, MA). The Cy3- and Cy5-labeled cDNA probes were mixed together and hybridized to a microarray slide. After overnight at 65 °C, the slide was washed twice with 2 $\times$  SSC containing 0.1% SDS for 5 min at 42 °C, once with 0.1 $\times$  SSC containing 0.1% SDS for 10 min at room temperature, and finally with 0.1 $\times$  SSC for 1 min at room temperature. Slide was dried by centrifugation at 650 rpm for 5 min. Hybridization images on the slide were scanned by Scanarray lite (Packard Bioscience, Boston, MA) and analyzed by GenePix Pro3.0 software (Axon Instrument, Union City, CA). Three independent experiments were performed, and the ratio of Cy3 and Cy5 signal intensity was calculated for each spot. These ratio was log<sub>2</sub>-transformed and normalized by subtracting the average of log<sub>2</sub> (Cy3/Cy5) values for internal control genes using Excel (Office 2000, Microsoft Corp.). For each gene, the mean values were then calculated and a difference of two-fold was applied to select up- or down-regulated genes by estrogen.

#### **4. Uterine sections and Laser capture microdissection**

Uterine horns were embedded in Tissue-Tek (Sakura Finetek, Torrance, CA) and frozen in liquid nitrogen. Cryosections (thickness, 6  $\mu\text{m}$ ) were cut and mounted onto clean glass slides. After the

sections were counterstained with Mayers hematoxylin, each population of uterine cells (luminal epithelial, muscle and stromal cells) was isolated from these sections using P.A.L.M. Robot-Microbeam version 4.0 (P.A.L.M. Microlaser Technologies AG, Bernried, Germany). For each cell, an average of 150 laser shots were transferred onto 0.5 ml tube cap and stored at -70 °C until utilized for total RNA extraction.

#### **5. Confirmation of microarray data with semiquantitative RT-PCR analysis**

Total RNA was extracted from whole uterine tissues or each population of uterine cells using TRIZOL reagent and purified using RNeasy total RNA isolation kit following the manufacturers instructions. One  $\mu\text{g}$  of RNA was reverse-transcribed at 42 °C for 60 min in 20  $\mu\text{l}$  reaction mixture consisting of oligo(dT)-adapter primer (Takara, Shiga, Japan) and AMV reverse transcriptase XL (Takara). The following PCR was performed in a total volume of 40  $\mu\text{l}$  with 2  $\mu\text{l}$  of the RT reaction mixture, 2  $\mu\text{l}$  25 mM MgCl<sub>2</sub>, 4  $\mu\text{l}$  10 $\times$  PCR buffer (100 mM Tris-HCl, 500 mM KCl, pH 8.3), 4  $\mu\text{l}$  2.5 mM dNTPs, 10 pmole forward and reverse primer and 1.25 U Taq polymerase (Takara). The sequences of the primers used were described in Table 3. An increasing number of cycles was tested to assess the best conditions of achieve linear amplification. The thermal cycling parameters consisted of 22~26 cycles of denaturing (94 °C, 30 sec); annealing (60 °C, 30 sec); and extension (72 °C, 30 sec). The PCR products were separated by electrophoresis on a 1.2% TBE agarose-ethidium bromide gels and visualized under UV light. The images were quantified by densitometric scanning followed by BioID image analysis software (Vilber-Lourmat, Mama La Vallee, Cedex, France) and gene expression was normalized against the density of the corresponding ribosomal protein L-7 (rpL7) PCR product as internal control.

**Table 1.** Genes up-regulated by estrogen in ovariectomized mouse uterus

Gene name	Fold change		Genbank No.
	6 H	12 H	
Cell cycle-related			
Nuclear factor I/C	-1.15	5.48	U57635
Protamine 2	1.21	3.11	X14004
Cyclin-dependent kinase inhibitor 1A (P21)	1.28	2.51	BC002043
Ubiquitin-activating enzyme E1C	1.31	2.22	AY029181
FBJ osteosarcoma oncogene	-1.05	2.22	BC029814
RAN, member RAS oncogene family	1.76	2.19	BC014829
Peroxisome proliferator activated receptor binding protein	1.04	2.07	AF000294
Serum-inducible kinase	-1.12	2.01	M96163
Immune-related			
Protein tyrosine phosphatase, receptor type, C	1.59	2.89	NM011210
Small chemokine (C-C motif) ligand 11	1.79	2.55	BC0027521
Proteasome subunit, beta type 5 (Psm5)	2.29	2.43	AF060091
Tumor necrosis factor (ligand) super family, member 13b	1.01	2.01	AF119383
Signal transduction			
Receptor (calcitonin) activity modifying protein 3 (Ramp3)	2.11	6.12	AF209907
Serine/arginine-rich protein specific kinase 2	1.26	5.96	BC020178
Chemokine orphan receptor 1	1.67	2.99	BC015254
Fibroblast growth factor 1	-1.19	2.94	U67610
Guanine nucleotide binding protein, alpha 12	1.49	2.68	M63659
Inositol polyphosphate-5-phosphatase, 145 kDa	-1.17	2.57	AF228679
Purinergic receptor P2Y, G-protein coupled 2	1.06	2.49	BC006613
Guanine nucleotide binding protein, alpha 11	1.93	2.23	M55411
Debrin-like	1.18	2.22	BC046430
Mitogen activated protein kinase 13	1.04	2.21	U81823
Endoglin	-1.04	2.14	X77952
MAD homolog 2 (Drosophila)	1.06	2.09	BC021342
Phosphatidylinositol 3-kinase, C2 domain containing, gamma	1.08	2.07	AB008792
Cell division cycle 42 homolog (S. cerevisiae)	1.19	5.79	L78075
Enzyme			
Branched chain ketoacid dehydrogenase E1	1.13	2.01	L16992
Branched chain aminotransferase 1, cytosolic	1.21	4.74	AK013888
Mevalonate (diphospho) decarboxylase	1.13	3.43	AJ309922
Mannosidase 1, beta	1.99	3.08	XM203179

**Table 1.** Continued

Gene name	Fold change		Genbank No.
	6 H	12 H	
Apoptosis-related			
Cytochrome c, somatic	1.87	2.79	BC034363
Sphingosine-1-phosphate phosphatase 1	-1.15	2.14	AF247177
Programmed cell death 6 interacting protein	1.33	2.13	C026823
Transcription			
Homeo box A5	1.73	9.63	X16840
Activating transcription factor 5 (AFT5)	1.44	4.03	BC010195
Zinc finger protein 1	-1.31	3.11	XM134378
Kruppel-like factor 4 (gut)	1.19	3.02	U20344
Paired-like homeodomain transcription factor 3	-1.19	2.86	AF005772
Williams-Beuren syndrome chromosome region 14	1.02	2.54	AF245479
General transcription factor II H, polypeptide 2	1.29	2.31	BC016231
CpG binding protein	-1.05	2.29	AY099096
Activating transcription factor 4 (ATF4)	-1.04	2.28	M94087
Paired related homeobox 1	-1.05	2.22	AK084387
Fibrillarin	1.16	2.21	Z22593
Ioquois related homeobox 3 (Drosophila)	1.14	2.17	Y15001
Tripartite motif protein 24	1.53	2.06	M64429
Activating transcription factor 1 (ATF1)	1.28	2.04	BC006871
Inhibitor of DNA binding 1 (Id-1)	1.23	2.01	M31885
Structure-related			
Small proline-rich protein 2A (Sprp2A)	1.95	75.86	BC010818
Small praline-rich protein 2H (Sprp2H)	1.27	12.07	AY158992
Procollagen, type I, alpha 1	-1.64	4.11	U03419
Small proline-rich protein A (SprpA)	-1.15	2.72	X91824
Procollagen, type III, alpha 1	-1.58	2.39	X52046
Others			
Pleckstrin homology, Sec7 and coiled/coil domain 3	1.25	7.04	NM011182
B-ell CLL/lymphoma 7B	1.52	4.77	AJ011145
Heat shock 70 kDa protein 4	-1.19	3.38	D85904
Immediately early response 2	1.77	3.11	BC002067
SEC61, gamma subunit ( <i>S. cerevisiae</i> )	2.85	2.52	U11027
Cystatin B	3.47	2.38	U59807
Eukaryotic translation initiation factor 2, subunit 2 (Eif2s2)	2.47	2.29	NM026030

**Table 2.** Genes down-regulated by estrogen in ovariectomized mouse uterus

Gene name	Fold change		Genbank No.
	6 H	12 H	
Cell cycle-related			
Nuclear factor I/A	-1.04	-3.79	Y07690
Epidermal growth factor receptor (EGFR)	-1.33	-2.44	BC023729
Checkpoint kinase 1 homolog (S. pombe)	-1.44	-2.28	AF16583
Topoisomerase (DNA) I	-2.11	-1.23	D10061
Immune-related			
Interleukin 15	ND	-7.74	U14332
Chemokine (C-X-C motif) ligand 12 (CXC12)	-3.19	-6.49	BC046827
Serine (or cysteine) proteinase inhibitor, Glade G, member 1	-1.54	-3.63	BC002026
Histocompatibility 2, class II antigen E beta	-1.23	-3.09	M36940
Fc receptor, IgG, alpha chain transporter	-1.19	-2.84	D37874
Immunoglobulin (CD79A) binding protein 1	-1.11	-2.75	XM196586
Complement component factor h	-2.19	-2.65	NM009888
Proteasome (prosome, macropain) subunit, beta type 9	-1.19	-2.16	BC032210
Chemokine (C-X-C motif) ligand 14 (CXC14)	-1.54	-2.13	AF192557
Chemokine (C-C motif) receptor 7	1.21	-2.04	L31580
Signal transduction			
Lymphoid blast crisis-like 1	-1.46	-9.73	U28495
Secreted frizzled-related sequence protein 2 (SFR2)	-1.41	-6.03	AF337040
Taste receptor, type 1, member 1	-1.29	-4.63	U36757
Coagulation factor II (thrombin) receptor	-1.04	-3.37	BC047086
Protein tyrosine phosphatase, receptor type, B	-1.30	-2.91	M84607
Platelet derived growth factor receptor, alpha polypeptide	1.23	-2.81	NM080428
F-box and WD-40 domain protein 7, archipelago homolog	-1.89	-2.63	D14340
Tight junction protein 1	-1.34	-2.57	AB028143
Endothelial differentiation, sphingolipid G-protein	-1.27	-2.41	U06924
Signal transducer and activator of transcription 1	-1.28	-2.29	AF194871
Interferon-inducible GTPase	-1.48	-2.25	U07617
Growth factor receptor bound protein 2	-1.29	-2.24	BC037696
Platelet-derived actor, C polypeptide	-1.25	-2.04	AF062484
Sorting nexin 12	-1.37	-2.03	BC014722
Enzyme			
UDP-Gal:betaGlcNAc beta 1,3-galactosyltransferase	-2.49	-11.97	BC046322
Putative phosphatase	1.24	-8.49	U96724
Thioether S-methyltransferase	-1.19	-3.52	M88694
Ubiquitin specific protease 21	-1.23	-2.99	BC021903
Cysteine dioxygenase 1, cytosolic	-1.72	-2.59	BC013638

**Table 2.** Continued

Gene name	Fold change		Genbank No.
	6 H	12 H	
Apoptosis-related			
Programmed cell death 5 (Pcd5)	-3.25	ND	AF161074
Tumor necrosis factor (ligand) superfamily, member 12	-1.55	-2.31	AF030100
Transcription			
Neurogenic differentiation 1 (NeuroD)	-2.28	ND	U28068
Transcription factor Dp1	-2.05	-3.15	X72310
Snail homolog 3 (Drosophila)	-1.62	-3.27	AF133714
Trichorhinophalangeal syndrome I (human)	-1.11	-2.89	AF346836
Y box protein 2	1.01	-2.63	AF073954
Histone deacetylase 5	-1.05	-2.41	AF207748
Ring inger protein 12	1.02	-2.33	AF069992
Eleven-nineteen lysine-rich leukemia gene	-1.14	-2.31	U80227
Lymphoid enhancer binding factor 1	-1.06	-2.29	X58636
Inhibitor of DNA binding 2 (Id-2)	-1.26	-2.26	M69293
Homeo box D9	-1.21	-2.13	BC019150
BTB and CNC homology 2 (Bach2)	-1.19	-2.08	D86604
WD40 protein Ciao 1	-1.01	-2.07	BC004089
Musculin (Msc)	-1.38	-2.04	AF087035
Structure-related			
Decorin	-4.75	-7.03	X53929
Fibromodulin	-1.37	-4.31	X94998
Tetranectin (plasminogen binding protein)	-1.51	-3.13	U08595
Syndecan 2	1.03	-2.13	U00674
Others			
Selenoprotein P, plasma, 1	-1.91	-7.99	X99807
Integrin alpha 9	-1.23	-4.18	AJ344342
RNA binding motif protein 5	-1.01	-3.09	AY309168
Integral membrane protein 2B (Imp2B)	-2.71	-2.91	U76253
Plexin B1	-1.02	-2.35	AB072381

## RESULTS

### 1. Global analysis of estrogen- regulated genes in ovariectomized mouse uterus

To analyze early and late estrogen-responsive

genes in the uterus, total uterine RNA was extracted at 6 h and 12 h after injection of estrogen to ovariectomized mice (OVX/estrogen treatment/ 6 h/12 h protocol) and the differential expression patterns were examined using TwinChip Mouse-7.4 K microarray. To confirm reproducibility, expe-


**Table 3.** Primer sequences for RT-PCR

Gene		Forward (F) and Reverse (S) primer sequences	Product size (bp)
Sprp2A	F	GTGTGGCCTGGGCCCTTGTCG	240
	R	GAGTCGGTGAGCTGGTGAG	
Cystatin B	F	CGCGCCATCTGCACAAT	225
	R	GGCTTGTTTTTCATGGGGGAG	
Ramp3	F	GGTGGTGTGGCGCAGCAAGC	427
	R	GCAGGGGTCAGGGTCAGGAC	
Psmb5	F	GGCTGGGGTGCAGCGGAT	350
	R	GGTAGATGGCTCGGCCGGG	
Eif2s2	F	CGTGATGGGGGTAAGGAGG	489
	R	CTAGAGCACAGGTTGGAG	
Id-1	F	GATCATGAAGGTCGCCAGTG	476
	R	TCCATCTGGTCTCAGTGC	
Decorin	F	CCCCTACCGATGCCAGTGTC	423
	R	GCTCCGTTTTCAATCCCAGA	
Imp2B	F	GTGGCGGTGGATTGCAAGGA	432
	R	GGGCGGCATACGATGGAAG	
SFR-2	F	TTGGCTTATACGTGCACT	295
	R	TATTTGAGGGCATCATGCAA	
Bach2	F	CCCATGTCACAAACCTATC	471
	R	TGCTCACCTGACACCGTTTCG	
CXC12	F	CGTGGGAGATGCAAGGGCAG	247
	R	GAGGAGAATGGGGATGAAGC	
Id-2	F	GTGACCAAGATGGAAATCCT	236
	R	TTTATTTAGCCACAGATAC	
rpL7	F	TCAATGGAGTAAGCCCAAAG	246
	R	CAAGAGACCGAGCAATCAAG	


periments were repeated independently three times and fold changes were calculated. The mean values of each spot intensities in the three experiments were calculated and plotted (Figure 1). Of the 7616 genes examined in OVX/estrogen treatment/6 h protocol, changes in mRNA expression were detected in 71 genes; 22 genes of these were activated and 49 genes were repressed (Table 1, 2). Only 0.9% of all genes were activated or re-

pressed more than two-fold and the remaining 99.1% of the genes revealed no significant differences in their expression by estrogen. However, 638 genes showed differential expression ratio more than or less than two-fold in OVX/estrogen treatment/12 h protocol; 351 genes of these were activated and 287 genes were repressed. About 8.4% of all genes were activated or repressed (Table 1, 2). Out of 638 activated or repressed genes, 433





**Figure 1.** MA-plots represent genes activated and repressed by estrogen in ovariectomized mouse uterus. MA-plot is used to represent the (R, G) data, where  $M = \log_2 R/G$  and  $A = \log_2 (R \times G)$ . (A) OVX/estrogen treatment/6 h protocol (B) OVX/estrogen treatment/12 h protocol. M, expression ratio; A, signal intensity; R, Red for Cy5; G, Green for Cy3.




**Figure 2.** Semiquantitative RT-PCR analysis confirming estrogen-regulated genes. Upper pannel, Expression patterns of estrogen-activated genes; Lower pannel, Expression patterns of estrogen-repressed genes. Values of each band were normalized to rpl7 for the same sample. Data were mean  $\pm$  SEM from three replicate experiments.

genes were listed as known gene in GenBank, and 205 genes were unknown. These genes were classified into eight functional categories as based on biological functions; cell cycle-related, immune-related, signal transduction, transcription-related, enzyme, structure-related, apoptosis-related, and others (including expressed sequence tag and un-

known genes).

## 2. Confirmation of estrogen-regulated genes

The expression patterns of several genes up- or down-regulated by estrogen were confirmed to verify the results of microarray analysis using semiquantitative RT-PCR. Several genes were ran-



**Figure 3.** Laser capture microdissection (LCM) procedure (A) Uterine sections stained with hematoxylin before LCM. LE, luminal epithelial cells; S, stromal cells; M, muscle cells. (B) Capture of luminal epithelial cells, (C) muscle cells and (D) stromal cells.

domly selected in different categories of genes; Sprp2A, Cystatin B, Ramp3, Psmb5, Eif2s2 and Id-1 (up-regulated), and Decorin, Imp2B, SFR2, Bach2, Id-2 and CXC12 (down-regulated). The results of semiquantitative RT-PCR analysis were shown in Figure 2.


### 3. Analysis of selected gene expression patterns in specific uterine cell types using LCM

To further analyze our findings we investigated gene expression of the selected up- or down-regulated genes in uterine specific cell types using laser capture microdissection technique (LCM). LCM allows the isolation of specific uterine cell types without contamination from other cell types. We isolated the population of specific uterine cell types; luminal epithelial, stromal and muscle cells (Figure 3). While Sprp2A, Eif2s2, and Psmb5

mRNAs were strongly activated in luminal epithelial cells, Ramp3 and Cystatin B were activated in both luminal epithelial and stromal cells. On the other hand, decorin mRNA was mainly repressed in muscle cells and the rest of down-regulated genes were repressed in stromal cells (Figure 4).

## DISCUSSIONS

Although estrogen plays an important role in various physiological events, the molecular mechanisms that are regulated by estrogen in the uterus remain largely unknown. Therefore, the identification of novel estrogen-regulated gene pathway is essential to understanding how estrogen regulates various uterine physiology, such as estrous cycle and implantation process. In the present study, we identified genes that are regulated



**Figure 4.** Expression patterns of estrogen-regulated genes in specific uterine cell types obtained with LCM. **(A)** Expression patterns of estrogen-activated genes. **(B)** Expression patterns of estrogen-repressed genes. C, oil-treated group; E 6H, post-estrogen 6 hr; E 12H, post-estrogen 12 hr; LE, luminal epithelial cells; S, stromal cells; M, muscle cells.

by estrogen in the ovariectomized mouse uterus exposed to  $17\beta$ -estradiol (E2) for 6 h and 12 h using cDNA microarray. Prior to use in cDNA microarray analysis, each uterine RNA sample via OVX/estrogen treatment/6 h/12 h protocol was validated by assaying for up-regulated (Asparagine synthetase and Lactoferrin) or down-regulated genes (Glutathione S-transferase and Secreted frizzled-related sequence protein 2) which were previously known to regulate by estrogen in mouse uterus.<sup>14,15</sup>

Our results showing that only 0.9% (71 genes) of all genes spotted in microarray slide were activated or repressed more than two-fold in OVX/estrogen treatment/6 h protocol suggest that direct or fast-responses by estrogen is not many in mRNA levels in uterus. However, 638 genes showed differential expression ratio more than or less than two-fold in OVX/estrogen treatment/12 h protocol. About 8.4% of all genes were activated

or repressed. It is possible that many indirect or nongenomic pathways affected the patterns of mRNA expression in latter protocol in addition to direct or genomic pathways. Recently, Watanabe et al. also performed the analysis of estrogen-responsive genes in ovariectomized mouse uterus.<sup>14</sup> They reported that generally 5% of all genes were activated or repressed more than three-fold by estrogen in OVX/estrogen treatment/6 h protocol; 616 genes were selected as estrogen-affected genes, of which 299 genes were activated and 317 genes were repressed. It is likely that this difference between their and our results is due to the species difference (they used C57/BL6/J strain) used in microarray analysis and the gene difference spotted in microarray.

Although our results to identify estrogen-regulated genes is quite different from that employed by Watanabe et al., two genes were shown the same expression pattern, chemokine orphan receptor 1 and receptor activity modifying protein 3 (Ramp3), which were slightly increased in OVX/estrogen treatment/6 h protocol and highly increased in OVX/estrogen treatment/12 h protocol. Calcitonin (CT), CT-gene-related peptide (CGRP), adrenomedullin (AM) and amylin are related hormones and neuropeptides. Ramp 1, 2 and 3, through interaction with calcitonin receptor (CTR), are required for the recognition of CGRP, AM and amylin by these receptor.<sup>16,17</sup> These relationship is elevated by estrogen and suggested a role for these genes in uterine estrogen response.

Activating transcription factor (ATF) 1, 4 and 5 were activated by estrogen in OVX/estrogen treatment/12 h protocol. It is known that ATF activate gene expression by binding as homo- or heterodimers to the cAMP response element in regulatory region of target genes. Although ATF family was not well characterized in uterine physiological events, such as estrous cycle and implantation process, its functions in other tissues suggest an

important role in cell proliferation and various signaling mechanisms.<sup>18,19</sup>

Small proline-rich protein 2A (Sprp2A) showed the highest level of up-regulation by estradiol in OVX/estrogen treatment/12 h protocol. This gene has been reported to express mainly in stratified squamous epithelial cells and control the differentiation of several cell types.<sup>20,21</sup> In our study, Sprp2A mRNA was dominantly activated in luminal epithelial cells (LE) compared to stromal and muscle cells using LCM techniques. This regulation suggested that the expression of Sprp2A in LE has a significance relationship between the epithelial cell function and developing embryos. Although the clarification of the function of Sprp2A and the significance of estrogen-responsiveness in relation to uterine physiological events are not identified, recent studies suggested that Sprp2A gene family may play a pivotal role in estrous cycle and implantation process.<sup>12,14</sup>

To the best of our knowledge this is the first study to profile the early- and lately-reponsive genes by estrogen involved in estrous and implantation process in mouse using cDNA microarray and LCM techniques. Our results indicate differential expression in spatiotemporal manner of genes regulated by estrogen and implicate various genes not previously known to express in ovariectomized mouse uterus. In conclusion, further analysis including to progesterone regulation, spatiotemporal expression, protein expression and function of these genes may be necessary to understand the exact mechanisms of estrogen effects in estrous and implantation process.

---

#### Acknowledgements

We thank Prof. Kang Hee-Gyoo for contributing to several experiments and helpful suggestions in Department of Biomedical Laboratory Science, Seoul Health College. We are also grateful to Lim Hee-Joung for excellent technical assistance of

laser capture microdissection.

#### REFERENCES

1. Green S, Walter P, Kumar V. Human oestrogen receptor cDNA: sequence, expression and homology to v-erb-A. *Nature* 1986; 320: 134-9.
2. Kuiper GG, Enmark E, Pelto-Huikko M, Nilsson S, Gustafsson JA. Cloning of a novel receptor expressed in rat prostate and ovary. *Proc Natl Acad Sci* 1996; 93: 5925-30.
3. Kousteni S, Bellido T, Plotkin LI. Nongenotropic, sex-nonspecific signaling through the estrogen or androgen receptors: dissociation from transcriptional activity. *Cell* 2001; 104: 719-30.
4. Angel N, Ana BR, Ouahiba L, Marjorie M, Esther F, Bernat S. Nongenomic actions of estrogens and xenoestrogens by binding at a plasma membrane receptor unrelated to estrogen receptor  $\alpha$  and estrogen receptor  $\beta$ . *Proc Natl Acad Sci* 2000; 97: 11603-8.
5. Ralf L, Martin W. Nongenomic actions of steroid hormones. *Mol cell Biol* 2003; 4: 46-56.
6. McMaster MT, Teng CT, Dey SK, Andre GK. Lactoferrin in the mouse uterus: Analysis of the preimplantation period and regulation by ovarian steroids. *Mol Endo* 1992; 5: 101-11.
7. Wang X, Das SK, Damm D, Klagsbrun M, Abraham JA, Dey SK. Differential regulation of heparin-binding epidermal growth factor-like growth factor in the adult ovariectomized mouse uterus by progesterone and estrogen. *Endocrinol* 1994; 135: 1264-71.
8. Das SK, Tsukamura H, Paria BC, Andrews GK, Dey SK. Differential regulation of epidermal growth factor receptor (EGF-R) gene and regulation of EGF-R bioactivity by progesterone and estrogen in the adult mouse uterus. *Endocrinol* 1994; 134: 971-81.
9. Diane MK, Sylvia CH, Kenneth SK, Richard PD. Activation of a uterine insulin-like growth factor I signaling pathway by clinical and environmental estrogens: requirement of estrogen receptor- $\alpha$ . *Endocrinol* 2000; 141: 3430-9.

10. Brannvall K, Korhonen L, Lindholm D. Estrogen-receptor-dependent regulation of neural stem cell proliferation and differentiation. *Mol Cell Neurosci* 2002; 21: 512-20.
11. Reese J, Das SK, Paria BC, Lim H, Song H, Matsumoto H, et al. Global gene expression analysis to identify molecular markers of uterine receptivity and embryo implantation. *J Biol Chem* 2001; 276: 44137-45.
12. Cheon YP, Li Q, Xu X, Demayo FJ, Bagchi IC, Bagchi MK. A genomic approach to identify novel progesterone receptor regulated pathways in the uterus during implantation. *Mol Endo* 2002; 16: 2853-71.
13. Green AR, Edwards RE, Greaves P, White INH. Comparison of the effect of oestradiol, tamoxifen and raloxifene on nerve growth factor- $\alpha$  expression in specific neonatal mouse uterine cell types using laser capture microdissection. *J Mol Endo* 2003; 30: 1-11.
14. Watanabe H, Suzuki A, Mizutani T, Khono S, Lubahn DB, Iguchi T. Genome-wide analysis of changes in early gene expression induced by estrogen. *Genes to Cells* 2002; 7: 497-507.
15. Das SK, Tan J, Johnson DC, Dey SK. Differential spatiotemporal regulation of lactoferrin and progesterone receptor genes in the mouse uterus by primary estrogen, catechol estrogen, and xenoestrogen. *Endocrinol* 1998; 139: 2905-15.
16. Fischer JA, Muff R, Born W. Functional relevance of G-protein-coupled-receptor-associated proteins, exemplified by receptor-activity-modifying proteins (RAMPs). *Biochem Soc Trans* 2002; 30: 455-60.
17. Gangula PR, Wimalawansa SJ, Yallampalli C. Pregnancy and sex steroid hormones enhance circulating calcitonin gene-related peptide concentrations in rats. *Hum Reprod* 2000; 5: 949-53.
18. Sabbah M, Courilleau D, Mester J, Redeuilh G. Estrogen induction of the cyclin D1 promoter: involvement of a cAMP response-like element. *Proc Natl Acad Sci* 1999; 96: 11217-22.
19. Bleckmann SC, Blendy JA, Rudolph D, Monahan AP, Schmid SG. Activating transcription factor 1 and CREB are important for cell survival during early mouse development. *Mol Cell Biol* 2002; 22: 1919-25.
20. Song HJ, Poly G, Darwiche N, Lichti U, Kuroki T, Kartasova T. Mouse *Spr2* genes: A clustered family of genes showing differential expression in epithelial tissues. *Genomics* 1999; 55: 28-42.
21. Fischer DF, Gibbs S, Putte P, Backendorf C. Interdependent transcription control elements regulate the expression of the *SPRP2A* gene during keratinocyte terminal differentiation. *Mol Cell Biol* 1996; 16: 5365-74.