

## Expression of Immunosuppression-Related Genes in Fetal Chorionic Villi Derived from Recurrent Spontaneous Abortion Patients

Ju-Mi Shin<sup>1,2</sup>, Jeong-Wook Kim<sup>4</sup>, Bumchae Choi<sup>4</sup>,  
Sook-Hwan Lee<sup>1,3</sup>, Kwang-Hyun Baek<sup>1,2,3\*</sup>

<sup>1</sup>Graduate School of Life Science and Biotechnology, <sup>2</sup>Cell and Gene Therapy Research Institute,  
<sup>3</sup>Pochon CHA University, Infertility Medical Center, CHA General Hospital, Seoul, Korea,  
<sup>4</sup>Department of Obstetrics and Gynecology, Creation and Love Women's Hospital, Kwangju, Korea

### 습관성 유산 환자의 융모막 조직에서의 면역억제유전자 발현 양상

포천중문 의과대학교 생명과학전문대학원<sup>1</sup>, 세포유전자치료연구소<sup>2</sup>,  
차병원 여성의학연구소<sup>3</sup>, 광주 시엘병원 산부인과<sup>4</sup>

신주미<sup>1,2</sup> 김정옥<sup>4</sup> 최범채<sup>4</sup> 이숙환<sup>1,3</sup> 백광현<sup>1,2,3\*</sup>

#### 연구목적:

연구재료 및 방법: 6 8  
(Normal N=6; RSA N=6).

2,3-dioxygenase (IDO) mucin1 (MUC1) Student's *t*-test , p<0.05  
placenta protein 14 (PP14), indoleamine  
GAPDH

결 과: 6 8 (PP14, IDO, MUC1)

결 론: 가 (PP14, IDO, MUC1)

**Key Words:** Chorionic villi, In situ hybridization, Pregnancy, Recurrent spontaneous abortion

During pregnancy, chorionic villi enlarge and blood vessels grow into them, forming highly vascularized structures, completely surrounding the chorion. These villi are important interface for

diffusion of nutrients, oxygen and wastes between mother and embryo. In these interface tissues, mother and fetus could consider to be a foreign each other, since a half of the fetal genome is

Tel: (02) 3468-3197, Fax: (02) 3468-3264, e-mail: baek@cha.ac.kr  
This study was supported by a grant of the Korea Health 21 R&D Project, Ministry of Health & Welfare, Republic of Korea (01-PJ10-PG6-01GN13-0002).

derived from the father. Recent investigations have been focused on immunological feto-maternal reaction that may cause recurrent spontaneous abortion (RSA).<sup>1</sup> RSA or habitual abortion, which is defined as the loss of three or more consecutive pregnancies before the 20th week of gestation, occurs in approximately 2~5% of pregnant women.<sup>2</sup> Why pregnancy loss containing RSA takes place at the molecular level remains unclear. It has been previously demonstrated that RSA can be caused by multiple problems associated with chromosomal abnormality, endocrine dysfunction, autoimmune disorders, and other unknown factors.<sup>3-5</sup>

One of immunosuppression-related molecules is glycodelin. It has been called by various names; placental protein 14 (PP14), chorionic  $\alpha$ 2-microglobulin (CAG-2), progesterone-associated endometrial protein (PEP) and pregnancy-associated  $\alpha$ 2-microglobulin ( $\alpha$ 2-PEG).<sup>6</sup> *PP14* has been isolated from human placenta and is expressed in other normal tissues including the epithelium of the fallopian tube, ovarian surface epithelium, uterine cervix, breast tissue, sweat glands, and bone marrow aspirates.<sup>7,8</sup> One of biological functions for PP14 is to inhibit early events in the T-cell receptor signaling pathway.<sup>6,9</sup> This indicates that PP14 has important roles as a possible immunosuppressive molecule capable of successful pregnancy and growth of the fetus.<sup>10,11</sup> Mucin1 (MUC1) is a cell-surface and secretory molecule of endometrial epithelium and shows the highest expression in the mid-luteal phase.<sup>12</sup> Mucins including MUC1 have various functions including protection against bacterial infections,<sup>13,14</sup> protection of proteins and cells from proteolysis, inhibition of cell attachment, promotion of cell attachment, and inhibition of immune cell function.<sup>15</sup> Indoleamine 2,3-dioxygenase (IDO) is an enzyme that catabolizes tryptophan, which induces the death of T-lymphocytes.<sup>16</sup> In murine pregnancy, IDO expression may suppress T-cell-mediated

induction of C3 deposition at the maternal-fetal interface that increases the risk of allogeneic fetal loss.<sup>17</sup> Taken all together, immunological regulation of these gene products is required for keeping normal pregnancy.

In this study, we compared the expression of three genes (*PP14*, *MUC1* and *IDO*) involved in immunosuppression in chorionic villi from RSA and normal patients to define the molecular regulation of immunological processes at the fetomaternal interface during pregnancy.

## MATERIALS AND METHODS

### 1. Tissue specimens

A total number of 12 fetal chorionic villi samples, 6 from RSA patients and 6 from patients with electively terminated pregnancy, were obtained from 12 pregnant women between 6 and 8 weeks of gestation (three samples for each gestation period). The chorionic villi from RSA patients were obtained from patients who had at least three times of unexplained miscarriages and from patients with no history of abortion, ectopic pregnancy, pre-term delivery, or stillbirth as controls. The average ages of RSA patients and fertile controls were  $34 \pm 6$  and  $32 \pm 4$  years, respectively. All specimens were obtained within 30 minute of collection, frozen, and stored at  $-80^\circ\text{C}$  until use. Informed written consent was obtained from all patients.

### 2. Immunocytochemical staining for fetal chorionic villi

Fetal chorionic villi from normal controls and RSA patients were immunocytochemically stained using antibodies for proliferating cell nuclear antigen (PCNA) to detect the proliferative activity, and hematoxylin and eosin (HE) to observe mitotic figures. Chorionic villi samples from normal and RSA patients were washed three times, 5 min

**Table 1.** Primers used for RT-PCR analysis. Primers of immunosuppression-related genes (*PPI4*, *IDO*, and *MUC1*) were made on the basis of published sequences in GenBank

Immunosuppression-related genes	Accession numbers	Nucleotide sequences	Size of PCR products
Placental protein 14	XM_005360	5' AAGTTGGCAGGGACCTGGCACTC 3' 5' ACGGCACGGCTCTCCATCTGTT 3'	422 bp
Indoleamine 2,3-dioxygenase	NM_002164	5' GCGCTGTTGGAAATAGCTTC 3' 5' CAGGACGTCAAAGCACTGAA 3'	119 bp
Mucin 1	NM_002456	5' AGACGTCAGCGTGAGTGATG 3' 5' CAGCTGCCCGTAGTTCTTTC 3'	171 bp
GAPDH	BC009081	5' ACCACAGTCCATGCCATCAC 3' 5' TCCACCACCCTGTTGCTGTA 3'	452 bp

each, in a 50 µl drop of PBS. This was followed by fixation at room temperature for 6 hrs in neutral buffered formalin. Then samples were washed and postfixed in 70% ethanol for 10 min. Villi samples were incubated with a 1 : 100 dilution of monoclonal anti-PCNA antibody (NCL-PCNA, Novocastra Lab., Newcastle upon Tyne, UK) for 18 hrs. They were washed three times for 5 min each time in Tris-buffer (pH 7.2). For the visualization of PCNA-labeled nuclei, villi were incubated for 10 min in the presence of biotinylated rabbit anti-mouse immunoglobulins at a dilution of 1 : 250. Also villi were incubated for 20 min in the presence streptavidin labeled horseradish peroxidase, and washed three times for 5 min in Tris-buffer. For the visualization of all nuclei, diaminobenzidine (DAB, Sigma Chemical Co., MO, USA) and Mayer's hematoxylin added and thereafter washed three times for 5 min. The washed villi were placed onto a coverslip and immediately mounted in xylene.

### 3. Total RNA preparation and cDNA synthesis

Total RNA was obtained from stored chorionic villi at -80 °C, using TRIzol reagent (Gibco BRL). cDNA was synthesized from total RNA according to the protocol of SuperScript Preamplification


system (Gibco BRL). After incubation of 3 µg total RNA with 0.5 µg oligo (dT)<sub>12-18</sub> primer at 70 °C for 10 min, the reaction was carried out in 5X first strand buffer, 10 mM DTT, and 0.5 mM dNTP containing final volume of 20 µl mixture. Mixed contents of the tube were incubated at 42 °C for 2 min. And then SUPERSRIPT™II (Gibco BRL) was added and incubated for 50 min at 42 °C.

### 4. Primers for reverse transcription-polymerase chain reaction (RT-PCR)

Primers of immunosuppression-related genes (*PPI4*, *IDO*, and *MUC1*) are made on the basis of published sequences in GenBank. (Accession numbers: XM\_005360, NM\_002164 and NM\_002456, respectively) The sequences for primers and their expected size of PCR products are listed in Table 1. The specific primers for *GAPDH* for a control are designed based on GeneBank accession number BC009081.

### 5. Semi-quantitative RT-PCR analysis

The relative expression level of genes isolated from RSA and normal chorionic villi samples was determined according to the expression level of *GAPDH*, a housekeeping gene. PCR condition is following; after boiling at 94 °C for 4 min, am-



**Figure 1.** Immunocytochemical staining for fetal chorionic villi from normal and RSA patients. Fetal chorionic villi were immunocytochemically stained with either an antibody for PCNA to detect the proliferating activity or HE to observe mitotic figures. Magnification X100. Bar=100  $\mu$ m.

plification was carried out for 30 cycles at 94 for 1 min, 55 for 1 min, and 72 for 2 min, the mixture was then cycled at 72 for 10 min to complete the elongation step and was finally stored at 4 . The density of each band was measured using a Gel-Doc image analyzer (Vilber Loumat, France). The band intensities of each gene were normalized with those of corresponding *GAPDH* used as a control.

### 6. Statistic analysis

All statistical analyses of the data, obtained from semi-quantitative RT-PCR for 3 different immunosuppression-related genes using 6 pairs of chorionic villi samples (Normal N=6; RSA N=6),


were performed using Student's *t*-test when normal and RSA groups were compared. Numerical data for the expression level of each gene are presented as mean  $\pm$  SD, and a p value less than 0.05 was considered statistically significant.

## RESULTS

The presence of live cells in two groups of chorionic villi derived from normal and RSA patients was confirmed by immunocytochemical analysis. Fetal chorionic villi from RSA patients and normal controls were immunocytochemically stained using hematoxylin and eosin (H/E) to observe mitotic figures and an antibody for prolife-

rating cell nuclear antigen (PCNA) to detect the proliferative activity. The analysis with hematoxylin and eosin (H/E) staining revealed that most of cells constituting chorionic villi in both groups are alive (Figures 1A and 1B). In addition, immunocytochemical analysis with anti-PCNA antibody (Figures 1C and 1D) showed that both groups of chorionic villi contain proliferative cells.


In order to investigate whether they are expressed abnormally in chorionic villi of RSA patients, we performed RT-PCR with primers generated for immunosuppression-related genes (*PP14*, *IDO* and




**Figure 2.** Semi-quantitative RT-PCR analysis in chorionic villi for immunosuppression-related genes. Immunosuppression-related genes tested in this study were *PP14*, *IDO* and *MUC1*. Samples were obtained from two groups, normal patients and RSA patients, at the gestation stages of 6 weeks and 8 weeks. (N; normal patients, RSA; RSA patients).

*MUC1*). Semi-quantitative RT-PCR analysis showed that these genes were less expressed in chorionic villi from the RSA group than in those from the therapeutic abortion group (Figures 2 and 3). It is noticeable that three genes showed different levels of expression between chorionic villi from normal controls and chorionic villi from RSA patients at both 6- and 8-weeks in gestation. To confirm the similar amount of RNA used for semi-quantitative RT-PCR analysis, a housekeeping gene *GAPDH* was used as a control (Figures 2 and 3). For semi-quantitative analyses for RT-PCR products, we first equalized band densities for *GAPDH* expression and then the expression level of each gene was compared between RSA and normal control patients that were gestational age matched (6 weeks and 8 weeks, respectively). The ratios were determined by dividing the band density of each gene by that of *GAPDH*. The intensity of the EtBr staining was analyzed by densitometry as described in MATERIALS AND METHODS, and plotted as a bar graph (Figure 3).

Statistical analysis revealed that the expression level of immunosuppression-related genes (*PP14*, *IDO* and *MUC1*) in fetal chorionic villi from normal pregnant women was significantly higher than that in fetal chorionic villi from pregnant women



**Figure 3.** Semi-quantitative RT-PCR analysis in chorionic villi for immunosuppression-related genes. Immunosuppression-related genes tested in this study were *PP14*, *IDO* and *MUC1*. The percent ratios were calculated for the expression of each gene in chorionic villi tissues from normal patients (open bar) and RSA patients (solid bar), respectively. The percent ratios were determined by dividing the band density of each gene by that of *GAPDH* followed by the multiplication of 100. All statistical analysis of the data was performed using Student's *t*-test ( $p < 0.05$ ).



**Figure 4.** Hypothesis for *in vivo* function of immunosuppression-related genes (*PP14*, *IDO* and *MUC1*). In general, they down-regulate immune reactions by inhibiting the proliferation of T-lymphocytes. The degree of immunosuppression activity in pregnant women is determined by immune cells and related cytokines.

with the RSA ( $p < 0.05$ ). The higher expression of three different immunosuppression-related genes occurred in six independent experiments (Normal  $N=6$ ; RSA  $N=6$ ). Taken all together, PP14, IDO, and MUC1 may have important roles as a possible immunosuppressive molecule capable of normal pregnancy and growth of fetus.

## DISCUSSION

The conceptus expresses both paternal and maternal antigens. Therefore, it is possible that the maternal immune system recognizes paternal gene products generated by the fetus as immunologically foreign, resulting in rejection.<sup>12,18,19</sup> However, the frequency of this rejection is rare due to the fact that uterine decidua secrete soluble proteins capable of inhibiting cell-mediated immune responses, protecting the conceptus from maternal immune rejection during pregnancy.<sup>20,21</sup> Approximately 25% of RSA patients has been shown the elevation of immune response to trophoblast and increased proliferation of inflammatory cells.<sup>22</sup> This leads to the possibility that aberrant regulation of

immunosuppression-related gene products may cause RSA or habitual abortion.

In this study, we investigated the possibility of aberrant expression level of immunosuppression-related genes (*PP14*, *IDO* and *MUC1*) in fetal chorionic villi of RSA patients. Molecular analysis showed that immunosuppression-related genes are expressed at a low level in chorionic villi from RSA patients than those from normal patients. This may result in high levels of immune reactions, following by RSA or habitual abortion. It has been demonstrated that PP14, IDO and MUC1 play a role in inhibiting the proliferation of T-lymphocytes.<sup>16,23,24</sup> In addition, down-regulation of the cellular immune response is dependent upon the suppression of T-helper (Th) 1 and T-cytotoxic (Tc) 1, which produce cytokines including IL-2, IFN- $\gamma$ , and TNF $\beta$ .<sup>25</sup> In the case of RSA patients, NK cells and cytolytic activity are increased due to the inability of inhibition for T-lymphocyte proliferation and of down-regulation for the cellular immune response.<sup>20,26</sup> Therefore, the degree of immunosuppression activity in pregnant women is determined by immune cells and related cytokines (Figure 4). This indicates that the appropriate regulation of immunosuppression-related gene expression is required for the early pregnancy state. Confirmation at the protein level, which was impossible with limited amounts of chorionic villi samples in our study, will help to verify their roles for normal pregnancy. This leads to the possibility that the identification of prognostic markers for RSA using immunosuppression-related gene products such as PP14, IDO and MUC1 will be helpful to provide some insights into the prognosis of pregnancies with a high risk of RSA and management of those pregnancies.

In addition to these genes, it is expected that there should be more genes involved in the process of establishing and maintaining pregnancy. Previous investigations have demonstrated that

RSA is a complicated problem associated with endocrine dysfunction, autoimmune disorders, chromosomal abnormalities, advanced maternal and paternal age, infectious processes, environmental toxins, and congenital or structural uterine anomalies,<sup>3,5,26-32</sup> indicating that a number of gene products may modulate these biological processes. Taken all together, the diagnosis of early pregnancy disorders including RSA depends on an accurate assessment of potential pathophysiological mechanisms and this will identify pregnancies with a high risk of miscarriage and enable to manage those pregnancies effectively.

---

#### Acknowledgements

We wish to thank the members of Cell and Gene Therapy Research Institute and Infertility Medical Center at CHA General Hospital, Pochon CHA University.

#### REFERENCES

- Hill JA, Choi BC. Maternal immunological aspects of pregnancy success and failure. *J Reprod Fertil* 2000; 55 (Suppl): 91-7.
- Coulam CB, Clark DA, Beer AE, Kutteh WH, Silver R, Kwak J, Stephenson M. Current clinical options for diagnosis and treatment of recurrent spontaneous abortion. *Am J Reprod Immunol* 1997; 38: 57-74.
- Bussen S, Sutterlin M, Steck T. Endocrine abnormalities during the follicular phase in women with recurrent spontaneous abortion. *Hum Reprod* 1999; 14: 18-20.
- Eblen AC, Gercel-Taylor C, Shields LB, Sanfilippo JS, Nakajima ST, Taylor DD. Alterations in humoral immune responses associated with recurrent pregnancy loss. *Fertil Steril* 2000; 73: 305-13.
- Lanasa MC, Higge WA, Kubik CJ, Ness RB, Harger J, Nagel R, Prosen T, Markobic N, Hoffman EP. A novel X chromosome-linked genetic cause of recurrent spontaneous abortion. *Am J Obstet Gynecol* 2001; 185: 563-8.
- Rachmilewitz J, Riely GJ, Tykocinski ML. Placental protein 14 functions as a direct T-cell inhibitor. *Cell Immunol* 1999; 191: 26-33.
- Dutta B, Mukhopadhyay D, Roy N, Das G, Karande AA. Cloning, expression, purification, and immunocharacterization of placental protein-14. *Protein Express Purifi* 1998; 14: 327-40.
- Connor JP, Brudney A, Ferrer K, Fazleabas AT. Glycodelin-A expression in the uterine cervix. *Gynec Oncol* 2000; 79: 216-9.
- Rachmilewitz J, Borovsky Z, Riely GJ, Miller R, Tykocinski ML. Negative regulation of T cell activation by placental protein 14 is mediated by the tyrosine phosphatase receptor CD45. *J Biol Chem* 2003; 278: 14059-65.
- Olajide F, Chard T. Biological and clinical significance of the endometrial protein PP14 in reproductive endocrinology. *Obstet Gynecol Surv* 1992; 47: 252-7.
- Lögberg L, Wester L. Immunocalins: a lipocalin subfamily that modulates immune and inflammatory responses. *Biochim Biophys Acta* 2000; 1482: 284-97.
- Hey NA, Li TC, Devine PL, Graham RA, Saravelos H, Aplin JD. MUC1 in secretory phase endometrium: expression in precisely dated biopsies and flushings from normal and recurrent miscarriage patients. *Hum Reprod* 1995; 10: 2655-62.
- Cohen MS, Black JR, Proctor RA, Sparling PF. Host defenses and the vaginal mucosa. *Scand J Urol Nephro* 1984; 86: 13-22.
- Lamblin G, Roussel P. Airway mucins and their role in defense against micro-organisms. *Respir Med* 1993; 87: 421-6.
- Lagow EM, DeSouza MD, Carson D. Mammalian reproductive tract mucins. *Hum Reprod* 1999; 5: 280-93.
- Sakurai K, Zou JP, Tschetter JR, Ward JM, Shearer GM. Effect of indoleamine 2,3-dioxygenase on experimental autoimmune encephalomyelitis. *Jor Ne*

- uroimmunol 2002; 129: 186-96.
17. Mellor AL, Keskin DB, Johnson T, Chandler P, Munn DH. Cells expressing indoleamine 2,3-dioxygenase inhibit T cell responses. *J Immunol* 2002; 168: 3771-6.
  18. Matsui S, Yoshimura N, Oka T. Characterization and analysis of soluble suppressor factor from early human decidual cells. *Transplantation* 1989; 47: 678-83.
  19. Dalton CF, Laird SM, Estdale SE, Saravelos HG, Li TC. Endometrial protein PP14 and CA-125 in recurrent miscarriage patients; correlation with pregnancy outcome. *Hum Reprod* 1998; 13: 3197-202.
  20. Laird SM, Tuckerman EM, Cork BA, Linjawi S, Blakemore AI, Li TC. A review of immune cells and molecules in women with recurrent miscarriage. *Hum Reprod Update* 2003; 9: 163-74.
  21. Kamarainen M, Leivo I, Koistinen R, Julkunen M, Karvonen U, Rutanen EM, Seppala M. Human ovary and ovarian tumors express glycodefin. A glycoprotein with immunosuppressive and contraceptive properties. *Am J Pathol* 1996; 148: 1435-43.
  22. Razaee A, Dabbagh A. T-helper (1) cytokine increase during early pregnancy in women with a history of recurrent spontaneous abortion. *Med Sci Monit* 2002; 8: 607-10.
  23. Mukhopadhyay D, Sunderesham S, Rao C, Karande AA. 2001. Placental protein 14 (PP14) induces apoptosis in T cells, but not in monocytes. *J Biol Chem* 2001; 276: 28268-73.
  24. Gimmi CD, Morrison BW, Mainprice BA, Gribben JG, Boussiotis VA, Freeman GJ, Park SY, Watanabe M, Gong J, Hayes DF, Kufe DW, Nadler LM. Breast cancer-associated antigen, DF3/MUC1, induced apoptosis activated human T cells. *Nat Med* 1996; 2: 1367-70.
  25. Michimata T, Sakai M, Miyazaki S, Ogasawara MS, Suzumori K, Aoki K, Nakata K, Saito S. Decrease of T-helper 2 and T-cytotoxic 2 cells at implantation sites occurs in unexplained spontaneous abortion with normal chromosomal content. *Hum Reprod* 2003; 18: 1523-8.
  26. Yamada H, Kato EH, Kobashi G, Ebina Y, Shimada S, Morikawa M, Sakuragi N, Fujimoto S. High NK cell activity in early pregnancy correlates with subsequent abortion with normal chromosomes in women with recurrent abortion. *Am J Reprod Immunol* 2001; 46: 132-6.
  27. Costa HD, de Moura MD, Feriani RA, Anceschi MI, Barbosa JE. Prevalence of anti cardiolipin antibody in habitual aborters. *Gynec Obstet Invest* 1993; 36: 221-5.
  28. Lim KJ, Odukoya OA, Li TC, Cooke ID. Cytokines and immuno-endocrine factors in recurrent miscarriage. *Hum Reprod Update* 1996; 2: 469-81.
  29. Coumans AB, Huijgens PC, Jakobs C, Schats R, de Vries JI, van Pampus MG, Dekker GA. Haemostatic and metabolic abnormalities in women with unexplained recurrent abortion. *Hum Reprod* 1999; 14: 211-4.
  30. Eblen AC, Gercel-Taylor C, Shields LBE, Sanfilippo JS, Nakajima ST, Taylor DD. Alterations in humoral immune responses associated with recurrent pregnancy loss. *Fertil Steril* 2000; 73: 305-13.
  31. Rai R, Backos M, Rushworth F, Regan L. Polycystic ovaries and recurrent miscarriage- a reappraisal. *Hum Reprod* 2000; 15: 612-5.
  32. Lee RM, Branch DW, Silver RM. Immunoglobulin A anti-beta2-glycoprotein antibodies in women who experience unexplained recurrent spontaneous abortion and unexplained fetal death. *Am J Obstet Gynecol* 2001; 185: 748-53.